

On quasi-Frobenius bimodules and corings

F. Castaño Iglesias *

Departamento de Estadística y Matemática Aplicada,

Universidad de Almería 04120, Almería, Spain

E-mail: fci@ual.es

1 Introduction

Frobenius bimodules are connected with Frobenius algebras and extensions. For instance, a ring extension $\varphi : R \rightarrow S$ is a Frobenius extension if and only if ${}_R S_S$ is a Frobenius bimodule [1]. Brzeziński and Gómez-Torrecillas studied in [4] certain properties of comatrix corings in relation to properties of bimodules. In particular they showed that the *comatrix coring* [6] induced by any Frobenius bimodule is a Frobenius coring. Recently, Castaño Iglesias and Năstăsescu [5] studied quasi-Frobenius extensions from the viewpoint of quasi-Frobenius functors. The ring extension $\varphi : R \rightarrow S$ is quasi-Frobenius (left and right extension) if and only if the restriction of scalars functor $\varphi_* : {}_S \mathcal{M} \rightarrow {}_R \mathcal{M}$ is a quasi-Frobenius functor (Remark 1.2).

Following this direction, we shall extend some results in [4] to the case when ${}_R M_S$ is a *quasi-Frobenius bimodule*. Using the notion of quasi-Frobenius functor and properties on quasi-Frobenius corings given in [5], we discuss the questions of when a quasi-Frobenius bimodule induce a quasi-Frobenius extension and a quasi-Frobenius coring.

Throughout this paper all rings will be assumed to have identity, and all modules to be unital. We use the standard module theory notation, for example a (R, S) -bimodule M is denoted by ${}_R M_S$, $\text{Hom}_R(-, -)$ denotes the Abelian group of R -module maps. The dual of a left R -module M is denoted by $({}_R M)^*$. Finally, ${}_R \mathcal{M}_S$ denotes the category of all (R, S) -bimodules. Let ${}_R M_S$ and ${}_R N_S$ be (R, S) -bimodules. M is said to be *similar* to N , abbreviated ${}_R M_S \sim {}_R N_S$, if there are $m, n \in \mathbb{N}$ and (R, S) -bimodules P and Q such that $M \oplus P \cong N^{(m)}$ and $N \oplus Q \cong M^{(n)}$ as (R, S) -bimodules (cf. [1]). It is easy to see that “ $\sim$ ” defines an equivalence relation on the class of (R, S) -bimodules. We start by recalling the definition of quasi-Frobenius functors between Grothendieck categories given in [5].

Let $\mathcal{A}$ be a Grothendieck category and consider an object X of $\mathcal{A}$. For any positive integer n , we denote by X^n the direct sum of n copies of X in the category $\mathcal{A}$. Given

*Research partially supported by Spanish Project (MTM2005–03227) from MCT.

additive and covariant functors $\mathbf{L}, \mathbf{R} : \mathcal{B} \rightarrow \mathcal{A}$ between Grothendieck categories, we say that $\mathbf{L}$ *divide* to $\mathbf{R}$, denoted by $\mathbf{L} \mid \mathbf{R}$, if for some positive integer n there are natural transformations

$$\mathbf{L}(X) \xrightarrow{\phi(X)} \mathbf{R}(X)^n \xrightarrow{\psi(X)} \mathbf{L}(X)$$

such that $\psi(X) \circ \phi(X) = 1_{\mathbf{L}(X)}$ for every $X \in \mathcal{B}$. $\mathbf{R} \mid \mathbf{L}$ is defined symmetrically. The functor $\mathbf{L}$ is said to be *similar* to $\mathbf{R}$, denoted by $\mathbf{L} \sim \mathbf{R}$, if $\mathbf{L} \mid \mathbf{R}$ and $\mathbf{R} \mid \mathbf{L}$.

1.1 Definition. The functor $\mathbf{F} : \mathcal{A} \rightarrow \mathcal{B}$ is said to be *quasi-Frobenius functor* if $\mathbf{F}$ has a left adjoint $\mathbf{L} : \mathcal{B} \rightarrow \mathcal{A}$ and also a right adjoint $\mathbf{R} : \mathcal{B} \rightarrow \mathcal{A}$ with $\mathbf{L}$ similar to $\mathbf{R}$.

1.2 Remark. Let $\varphi : R \rightarrow S$ be a ring extension. Associated to φ we have the restriction of scalars functor $\varphi_* : {}_S\mathcal{M} \rightarrow {}_R\mathcal{M}$, the induction functor $S \otimes_R - : {}_R\mathcal{M} \rightarrow {}_S\mathcal{M}$ and the coinduction functor $\text{Hom}_R({}_R S_S, -) : {}_R\mathcal{M} \rightarrow {}_S\mathcal{M}$. It is well known that φ_* is right adjoint to $S \otimes_R -$ and left adjoint to $\text{Hom}_R({}_R S_S, -)$. Then φ_* is a *quasi-Frobenius functor* if the functor $S \otimes_R -$ is similar to $\text{Hom}_R({}_R S_S, -)$. This is equivalent to assuming ${}_R S$ finitely generated projective and ${}_S S_R \sim \text{Hom}_R({}_R S_S, R)$. On the other hand, recall of [7] that a ring homomorphism $\varphi : R \rightarrow S$ is called *left quasi-Frobenius extension* if ${}_R S$ is finitely generated and projective and ${}_S S_R \mid \text{Hom}_R({}_R S_S, R)$. Equivalently, S_R and ${}_R S$ are finitely generated projective and $\text{Hom}_R(S_R, R) \mid {}_R S_S$.

Similarly, φ is called *right quasi-Frobenius extension* if S_R is finitely generated projective and ${}_R S_S \mid \text{Hom}_R(S_R, R)$ (or $\text{Hom}_R({}_R S_S, R) \mid {}_S S_R$). Then φ is a *quasi-Frobenius extension* (left and right extension) if ${}_R S$ is a finitely generated projective and ${}_S S_R \sim \text{Hom}_R({}_R S_S, R)$ (or ${}_R S_S \sim \text{Hom}_R(S_R, R)$). Therefore, the ring extension φ is quasi-Frobenius if and only if $\varphi_* : {}_S\mathcal{M} \rightarrow {}_R\mathcal{M}$ is a quasi-Frobenius functor.

2 Quasi-Frobenius bimodules

In this section we generalize the notion of Frobenius bimodule to *quasi-Frobenius bimodule*. Recall of [1] that a (R, S) -bimodule M is called Frobenius bimodule if ${}_R M$ and M_S are finitely generated and projective modules and $({}_R M)^* \cong (M_S)^*$ as (S, R) -bimodules. We start with the following definition.

2.1 Definition. An (R, S) -bimodule M is said to be *quasi-Frobenius bimodule*, if both ${}_R M$ and M_S are finitely generated projective and $({}_R M)^* \sim (M_S)^*$ as (S, R) -bimodules.

For any ring extension $\varphi : R \rightarrow S$, it is easy to see that φ is a quasi-Frobenius extension if and only if the natural bimodule ${}_R S_S$ is a quasi-Frobenius bimodule. So quasi-Frobenius bimodules generalize quasi-Frobenius extension.

2.2 Remark. It is obvious that the class of quasi-Frobenius bimodule contains Frobenius bimodules. In general, a quasi-Frobenius bimodule need not be a Frobenius (an example of finite-dimensional quasi-Frobenius algebra which is not a Frobenius is gives in [8]).

The following shows some relations between quasi-Frobenius functors and bimodules. If ${}_R M_S$ and ${}_S N_T$ are bimodules, then $M \otimes_S N$ receives the natural (R, T) -bimodule structure indicated by $r(m \otimes n)t = rm \otimes nt$.

2.3 Proposition. *Suppose ${}_R M_S$ and ${}_S N_T$ are quasi-Frobenius bimodules. Then ${}_R(M \otimes_S N)_T$ is a quasi-Frobenius (R, T) -bimodule.*

Proof. First, note that $({}_R(M \otimes_S N))^*$ and $((M \otimes_S N)_T)^*$ are finitely generated and projective modules, since ${}_R M$, M_S , ${}_S N$ and N_T are finitely generated projective modules. Next we apply $\text{Hom}_S({}_S N, -)$ to $({}_R M)^* \sim (M_S)^*$ to obtain:

$$\text{Hom}_S({}_S N, ({}_R M)^*) \sim \text{Hom}_S({}_S N, (M_S)^*)$$

Equivalently, $({}_R(M \otimes_S N))^* \sim ((M \otimes_S N)_S)^*$. Similarly, applying $\text{Hom}_S(M_S, -)$ to $({}_S N)^* \sim (N_T)^*$, it follows that $((M \otimes_S N)_S)^* \sim ((M \otimes_S N)_T)^*$. Thus $({}_R(M \otimes_S N))^* \sim ((M \otimes_S N)_T)^*$, whence $M \otimes_S N$ is a quasi-Frobenius (R, T) -bimodule. $\square$

2.4 Corollary. *If $\varphi : R \rightarrow S$ and $\psi : S \rightarrow T$ are quasi-Frobenius ring extensions, then ${}_R T_T$ is a quasi-Frobenius bimodule.*

Proof. The corollary follows from letting $M = {}_R S_S$ and $N = {}_S T_T$ in the proposition. $\square$

Let M be any (R, S) -bimodule. It is well-known that the functor $M \otimes_S - : {}_S \mathcal{M} \rightarrow {}_R \mathcal{M}$ has a right adjoint $\text{Hom}_R({}_R M, -) : {}_R \mathcal{M} \rightarrow {}_S \mathcal{M}$. If in addition, M_S is a finitely generated and projective module, then the functor $M \otimes_S -$ has also a left adjoint $(M_S)^* \otimes_R - : {}_R \mathcal{M} \rightarrow {}_S \mathcal{M}$. When this is the case, $M \otimes_S -$ is a quasi-Frobenius functor if the functors $(M_S)^* \otimes_R -$ and $\text{Hom}_R({}_R M, -)$ are similar. This leads to a proposition needed later:

2.5 Proposition. *For any (R, S) -bimodule M the following assertions are equivalent.*

- (a) ${}_R M_S$ is a quasi-Frobenius bimodule;
- (b) $M \otimes_S -$ is a quasi-Frobenius functor.

Proof. (a) $\Rightarrow$ (b). Suppose ${}_R M$ and M_S are finitely generated projective modules and $(M_S)^* \sim ({}_R M)^*$. Then by [5, Lemma 2.1], $(M_S)^* \otimes_R - \sim ({}_R M)^* \otimes_R -$. We note that $({}_R M)^* \otimes_R - \cong \text{Hom}_R({}_R M, -)$, since ${}_R M$ is a finitely generated projective. Therefore, $(M_S)^* \otimes_R -$ and $\text{Hom}_R({}_R M, -)$ are similar, which implies that $M \otimes_S -$ is a quasi-Frobenius functor.

(b) $\Rightarrow$ (a). Assume that $M \otimes_S -$ is a quasi-Frobenius functor. By [5, Theorem 2.2], ${}_R M_S$ is a finitely generated projective on both sides with $(M_S)^* \otimes_R - \sim \text{Hom}_R({}_R M, -)$. So $(M_S)^* \otimes_R R \sim \text{Hom}_R({}_R M, R)$, whence $(M_S)^* \sim ({}_R M)^*$. Consequently, ${}_R M_S$ is a quasi-Frobenius bimodule. $\square$

As a immediate consequence of the proposition above with $M = {}_R S_S$ and Remark 1.2, we have

2.6 Corollary. *Let $\varphi : R \rightarrow S$ be a ring extension. Are equivalent the following assertions*

- (i) φ is a quasi-Frobenius extension.
- (ii) ${}_R S_S$ is a quasi-Frobenius bimodule.
- (iii) $\varphi_* : {}_S \mathcal{M} \rightarrow {}_R \mathcal{M}$ is a quasi-Frobenius functor.

Consider now an (R, S) -bimodule M and let $T = \text{End}_S(M_S)$ be the endomorphism ring of M_S . Then there is a ring extension $\overline{\varphi} : R \rightarrow T$ defined by $r \mapsto \overline{\varphi}_r(m) = rm$. Note too the natural bimodule ${}_T M_S$ given by $tms = t(m)s$. The next theorem generalize the classical endomorphism ring theorem to the case of quasi-Frobenius bimodules.

2.7 Theorem. *If ${}_R M_S$ is a quasi-Frobenius bimodule, then $T = \text{End}_S(M_S)$ is a quasi-Frobenius extension of R .*

Proof. By Definition 2.1, it suffices to prove that ${}_R T_T$ is a quasi-Frobenius bimodule. Applying $M \otimes_S -$ to $(M_S)^* \sim ({}_R M)^*$, we obtain $M \otimes_S (M_S)^* \sim M \otimes_S ({}_R M)^*$. This establishes that $T = \text{End}_S(M_S) \cong M \otimes_S (M_S)^* \sim M \otimes_S ({}_R M)^* = ({}_R T)^*$. On the other hand, note that ${}_R T$ is finitely generated projective, since M_S and (M_S) are finitely generated projective modules. So ${}_R T_T$ is a quasi-Frobenius bimodule. $\square$

A converse of this theorem is give below where (1) is a *condition Willard's*, which is satisfied by a generator module [10].

2.8 Proposition. *Let R and S be rings and M a (R, S) -bimodule such that ${}_R M$ and M_S are both finitely generated projective modules. Suppose $T = \text{End}_S(M_S)$ is a quasi-Frobenius extension of R , and*

$$\text{Hom}_T({}_T M_{S,T} {}_T R) \cong \text{Hom}_S({}_R M_S, S_S). \quad (1)$$

Then ${}_R M_S$ is a quasi-Frobenius bimodule.

Proof. In the computation below, we apply the Hom-Tensor Relation, the necessary condition $({}_R T)^* \sim (T_T)^*$ for a quasi-Frobenius extension $R \rightarrow T$ and condition Willard $({}_T M)^* \cong (M_S)^*$ in that order:

$$\begin{aligned} ({}_R M)^* = \text{Hom}_R({}_R M, R) &= \text{Hom}_R(T \otimes_T M, R) \\ &\cong \text{Hom}_T({}_T M, \text{Hom}_R({}_R T, R)) \\ &= \text{Hom}_T({}_T M, ({}_R T)^*) \\ &\sim \text{Hom}_T({}_T M, (T_T)^*) \\ &\cong \text{Hom}_T({}_T M, T) = ({}_T M)^* \cong (M_S)^* \end{aligned}$$

Now the proposition follows, since ${}_R M$ and M_S are both finitely generated projective modules. $\square$

3 Application to corings

Let A be an associative and unitary algebra over a commutative ring (with unit) k . We recall from [9] that an A -coring $\mathfrak{C}$ consists of an A -bimodule $\mathfrak{C}$ with two A -bimodule maps

$$\Delta : \mathfrak{C} \rightarrow \mathfrak{C} \otimes_A \mathfrak{C}, \quad \epsilon : \mathfrak{C} \rightarrow A$$

such that $(\mathfrak{C} \otimes_A \Delta) \circ \Delta = (\Delta \otimes_A \mathfrak{C}) \circ \Delta$ and $(\mathfrak{C} \otimes_A \epsilon) \circ \Delta = (\epsilon \otimes_A \mathfrak{C}) \circ \Delta = 1_{\mathfrak{C}}$. A right $\mathfrak{C}$ -comodule is a pair (M, ρ_M) consisting of a right A -module M and an A -linear map $\rho_M : M \rightarrow M \otimes_A \mathfrak{C}$ satisfying $(M \otimes_A \Delta) \circ \rho_M = (\rho_M \otimes_A \mathfrak{C}) \circ \rho_M$ and $(M \otimes_A \epsilon) \circ \rho_M = 1_M$. The right $\mathfrak{C}$ -comodules together with their morphisms form the additive category $\mathcal{M}^{\mathfrak{C}}$. If ${}_A \mathfrak{C}$ is flat, then $\mathcal{M}^{\mathfrak{C}}$ is a Grothendieck category.

The forgetful functor $\mathbf{U} : \mathcal{M}^{\mathfrak{C}} \rightarrow \mathcal{M}_A$ has the right adjoint $- \otimes_A \mathfrak{C}$ (see [3, Lemma 3.1]). When ${}_A \mathfrak{C}$ is finitely generated and projective, the functor $\text{Hom}_A(\mathfrak{C}, -) : \mathcal{M}_A \rightarrow \mathcal{M}^{\mathfrak{C}}$ is a left adjoint to $\mathbf{U}$. In this case, $\mathbf{U}$ is a quasi-Frobenius functor if $- \otimes_A \mathfrak{C}$ and $\text{Hom}_A(\mathfrak{C}, -)$ are similar.

3.1 Definition. An A -coring $\mathfrak{C}$ is called *quasi-Frobenius coring* provided the forgetful functor $\mathbf{U} : \mathcal{M}^{\mathfrak{C}} \rightarrow \mathcal{M}_A$ is a quasi-Frobenius functor.

A characterization of such corings is given in [5, Theorem 5.5]. In particular an A -coring $\mathfrak{C}$ is a quasi-Frobenius coring if and only if ${}_A \mathfrak{C}$ is finitely generated projective and $\mathfrak{C} \sim E$ as (A, E) -bimodules where E is the opposite algebra of $\mathfrak{C}^*$.

Associated to any (R, S) -bimodule M we can consider the *comatrix S -coring* $(M_S)^* \otimes_R M$ provided M_S is a finitely generated projective [6]. As for Frobenius bimodules we obtain the following theorem, which generalizes [4, Theorem 3.7].

3.2 Theorem. *Let ${}_R M_S$ be a bimodule and $T = \text{End}_S(M_S)$ be the endomorphism ring. Then*

- (a) *If ${}_R M_S$ is a quasi-Frobenius bimodule, then $(M_S)^* \otimes_R M$ is a quasi-Frobenius S -coring.*
- (b) *If $(M_S)^* \otimes_R M$ is a quasi-Frobenius S -coring, then $T \otimes_R T$ is a quasi-Frobenius T -coring.*

Proof. (a). Suppose that ${}_R M_S$ is a quasi-Frobenius bimodule. Then the functors $(M_S)^* \otimes_R -$ and $\text{Hom}_R({}_R M, -)$ are similar by Proposition 2.5. So $(M_S)^* \otimes_R M \sim \text{Hom}_R({}_R M, M)$. From [6, Proposition 2.1], $\text{Hom}_R({}_R M, M) \cong E$ where E is the opposite algebra of $((M_S)^* \otimes_R M)^*$. Thus $(M_S)^* \otimes_R M \sim E$. Now [5, Theorem 4.4] implies that $(M_S)^* \otimes_R M$ is a quasi-Frobenius coring, since $(M_S)^* \otimes_R M$ is a finitely generated projective left S -module by assumptions.

(b). Suppose that $(M_S)^* \otimes_R M$ is a quasi-Frobenius S -coring. Then $(M_S)^* \otimes_R M \sim \text{Hom}_R({}_R M, M)$. Applying first the functor $M \otimes_S -$ and later $- \otimes_S (M_S)^*$, we obtain

$$M \otimes_S (M_S)^* \otimes_R M \otimes_S (M_S)^* \sim M \otimes_S \text{Hom}_R({}_R M, M) \otimes_S (M_S)^*$$

This is equivalent to

$$T \otimes_R T \sim \text{Hom}_R(T, M \otimes_S (M_S)^*) \cong \text{Hom}_R(T, T) \cong \overline{E}$$


where $\overline{E}$ is the opposite algebra of $((_R T)^* \otimes_R T)^*$. Since $(_R T)^* \otimes_R T \cong T \otimes_R T$, we note that [5, Theorem 4.4] finish the proof. $\square$

Following Sweedler [9], given a ring extension $\rho : R \rightarrow S$ one can view $\mathfrak{C} = S \otimes_R S$ as an S -coring. $\mathfrak{C}$ is known as *Sweedler's coring* associated to ρ . As an immediate consequence of Theorem 3.2 we obtain the following corollary.

3.3 Corollary. *If ${}_R S_S$ is a quasi-Frobenius bimodule, then the Sweedler's coring $S \otimes_R S$ is a quasi-Frobenius coring.*

3.4 Remark. A similar proof to Brzeziński [2, Theorem 2.7] show that $R \rightarrow S$ is a quasi-Frobenius extension whenever ${}_R S$ is faithfully flat and $S \otimes_R S$ is a quasi-Frobenius coring.

3.5 Remark. Let ${}_R M_S$ be a bimodule and $T = \text{End}_S(M_S)$ be the ring endomorphism. Then the following diagram summarise the main results of this paper.



References

- [1] F.W. Anderson and K.R. Fuller, *Rings and Categories of Modules*, GTM 13 Springer-Verlag, Berlin, 1974.
- [2] T. Brzeziński, *Towers of Corings*, Commun. Algebra **31**(4) (2003), 2015-2026.
- [3] T. Brzeziński, *The structure of corings. Induction functors, Maschke-type theorem, and Frobenius and Galois-type properties*, Algebras and Representation Theory **5** (2002), 389-410.
- [4] T. Brzeziński and J. Gómez-Torrecillas, *On Comatrix Corings and Bimodules*, K-Theory **29** (2003), 101-115.
- [5] F. Castaño Iglesias and C. Năstăsescu, *Quasi-Frobenius functors with applications to corings*, arXiv:math.RA/0612662, 2006.

- [6] L. El Kaoutit and J. Gómez-Torrecillas, *Comatrix corings: Galois corings, descent theory, and a structure theorem for cosemisimple corings*, Math. Z. **30** (2002), 2405-2426.
- [7] B. Müller, *Quasi-Frobenius-Erweiterungen*, Math. Z. **85** (1964), 345–368.
- [8] T. Nakayama, *On Frobenius algebras, I*, Ann. of Math. **40** (1939), 611-633.
- [9] M. E. Sweedler. *The predual theorem to the Jacobson-Bourbaki theorem*, Trans. Amer. Math. Soc. **213** (1975), 391-406.
- [10] E.R. Willard, *Properties of projective generators*, Math. Ann. **158** (1965), 352–364.